arXiv:1101.4806v1 [math.NT] 25 Jan 2011

STERN'S TYPE CONGRUENCES FOR GENERALIZED BERNOULLI NUMBERS

HAO PAN AND YONG ZHANG

1. Introduction

The Bernoulli numbers B_n and the Euler numbers E_n are given by

$$\sum_{n=0}^{\infty}\frac{B_n}{n!}t^n=\frac{t}{e^t-1},\qquad\qquad \sum_{n=0}^{\infty}\frac{E_n}{n!}t^n=\frac{2}{e^t+e^{-t}}.$$

The well-known Kummer theorem asserts that for a prime $p\geq 5$ and even $k,l>0$,

$$\frac{B_k}{k}\equiv\frac{B_l}{l}\ (\mathrm{mod}\ p)$$

whenever $k\equiv l\ (\mathrm{mod}\ p-1)$ and $p-1\nmid k$. In fact, Kummer also proved a similar result for the Euler numbers:

$$E_k\equiv E_l\ (\mathrm{mod}\ p),$$

provided that p is an odd prime, $k,l\geq 0$ are even and $k\equiv l\ (\mathrm{mod}\ p-1)$.

However, for E_n, the case $p=2$ is a little special. A result of Stern says that for even $k,l\geq 0$,

$$E_k\equiv E_l\ (\mathrm{mod}\ 2^n)$$

if and only if $k\equiv l\ (\mathrm{mod}\ 2^n)$. Clearly Stern's result can be restated as the congruence

$$E_{k+2^nq}\equiv E_k+2^n\ (\mathrm{mod}\ 2^{n+1}) \tag{1.1}$$

for even $k\geq 0$ and odd $q\geq 1$. Recently, Z.-H. Sun [1] extended the Stern congruence and showed that

$$E_{k+2^mq}\equiv E_k+2^mq\ (\mathrm{mod}\ 2^{m+2})$$

for $m\geq 2$ and

$$E_{k+2^mq}\equiv E_k+5\cdot 2^mq\ (\mathrm{mod}\ 2^{m+2})$$

for $m\geq 3$. For the history and a proof of Stern's congruence, the reader may refer to Wagstaff's survey [3].

For a Dirichlet character χ modulo m, define the generalized Bernoulli numbers $B_{n,\chi}$ by

$$\sum_{n=0}^{\infty}\frac{B_{n,\chi}}{n!}t^n=\sum_{a=1}^{m}\chi(a)\frac{te^{at}}{e^{mt}-1}.$$

Note that for $m>1$

$$\sum_{n=0}^{\infty}\frac{B_{n,\chi}}{n!}(-t)^n=\frac{1}{m}\sum_{a=1}^{m-1}\chi(a)\frac{(-t)e^{-at}}{e^{-mt}-1}=\frac{1}{m}\sum_{a=1}^{m-1}\chi(a)\frac{te^{(m-a)t}}{e^{mt}-1}=\frac{\chi(-1)}{m}\sum_{a=1}^{m}\chi(a)\frac{te^{at}}{e^{mt}-1}.$$

So $B_{n,\chi}=0$ if n and χ have the opposite parity. Let

$$\beta_{n,\chi}=\frac{B_{n,\chi}}{n}$$

for $n\geq 1$. It is easy to see that

$$E_k=-2\beta_{k+1,\chi_4},$$

where χ_4 is the unique non-trivial character modulo 4.

It is natural to ask whether there exist some similar congruences for the general $\beta_{n,\chi}$.

Theorem 1.1. *Suppose that $m \geq 3$ and χ be a primitive character χ modulo 2^m. Let $q \geq 1$ be odd and $n \geq 1$. If $k \geq 1$ has the same parity as χ, then $(1-\chi(5))\beta_{k,\chi}$ is an algebraic integer, and*

$$(1-\chi(5))(\beta_{k+2^n q,\chi} - \beta_{k,\chi}) \equiv \chi(5)2^{n+2}\beta_{d,\chi} \pmod{2^{n+2}}, \tag{1.2}$$

where $d \in \{1,2\}$ satisfying that $k \equiv d \pmod 2$. In particular, we have

$$(1-\chi(5))\beta_{k+2^n q,\chi} \equiv (1-\chi(5))\beta_{k,\chi} \pmod{2^{n+1}}. \tag{1.3}$$

In fact, we can get the congruences modulo the higher powers of 2.

Theorem 1.2. *Under the assumptions of Theorem 1.1, letting $d \in \{1,2,3,4\}$ with $k \equiv d \pmod 4$, if χ is even, then for each even $k \geq 2$,*

$$\begin{aligned}&(1-\chi(5))(\beta_{k+2^n q,\chi} - \beta_{k,\chi}) - 2^{n+2}(q+2)\beta_{d,\chi}\\ \equiv&2^{n-1}\big((1-\chi(5)5^2)(1+q)\beta_{2,\chi} + (1-\chi(5)5)(3q-5)\beta_{1,\chi\chi_4}\big) \pmod{2^{n+3}}.\end{aligned} \tag{1.4}$$

If χ is odd, then for each odd $k \geq 1$,

$$\begin{aligned}&(1-\chi(5))(\beta_{k+2^n q,\chi} - \beta_{k,\chi}) - \frac{5-\chi(5)5}{1-\chi(5)5}\cdot 2^{n+2}(q+2)\beta_{d,\chi}\\ \equiv&2^{n-1}\big((1-\chi(5)5^2)(1-q)\beta_{2,\chi\chi_4} + 3(1-\chi(5)5)(1-q)\beta_{1,\chi}\big) \pmod{2^{n+3}}.\end{aligned} \tag{1.5}$$

Similarly, for odd prime p, we have

Theorem 1.3. *Let p be an odd prime and χ be a primitive character χ modulo p^m, where $m \geq 1$. Let n and q be positive integers with $p \nmid q$. Suppose that $k \geq 1$ and k has the same parity as χ.*

(i) *Suppose that $\chi(a)a^k - 1$ is prime to p for an integer a with $p \nmid a$. Then $\beta_{k+p^n(p-1)q,\chi}$ is an algebraic integer, and*

$$\beta_{k+p^n(p-1)q,\chi} \equiv \beta_{k,\chi} \pmod{p^{n+1}}. \tag{1.6}$$

(ii) *Suppose that $m \geq 2$. Then $(1-\bar{\chi}(p+1))\beta_{k+p^n(p-1)q,\chi}$ is an algebraic integer, and*

$$(1-\bar{\chi}(p+1))(\beta_{k+p^n(p-1)q,\chi} - \beta_{k,\chi}) \equiv p^{n+1}q\beta_{d,\chi} \pmod{p^{n+1}}, \tag{1.7}$$

where $d \in \{1,2,\ldots,p-1\}$ with $k \equiv d \pmod{p-1}$.

However, the congruences modulo p^{n+2} for $\beta_{k+p^n(p-1)q,\chi} - \beta_{k,\chi}$ is a little complicated.

Theorem 1.4. *Under the assumptions of Theorem 1.3, we have the following results:*

(i) *Suppose that $\chi(a)a^k - 1$ is prime to p for an integer a with $p \nmid a$. Then*

$$\begin{aligned}&\beta_{k+p^n(p-1)q,\chi} - \beta_{k,\chi} - \frac{p^n(a^{p-1}-1)}{1-\chi(a)a^d}\cdot\chi(a)a^d q\beta_{d,\chi}\\ \equiv&q\sum_{\lambda \bmod p}\frac{1-\chi\lambda(a)a^d}{1-\chi(a)a^d}\beta_{d,\chi\lambda}\sum_{t=1}^{p-1}\bar{\lambda}(t)t^{d-1}(p^{n-1}(t^{p-1}-1)+p^n t^{p-2}) \pmod{p^{n+2}},\end{aligned} \tag{1.8}$$

where $d \in \{1,2,\ldots,p-1\}$ with $k \equiv d \pmod{p-1}$.

(ii) *Suppose that* $m \geq 2$ *and* $p \geq 5$. *Then*

$$(1-\bar{\chi}(p+1))(\beta_{k+p^n(p-1)q,\chi}-\beta_{k,\chi})-\left(1+\left(d-\frac{3}{2}\right)p-\frac{dp}{1-\bar{\chi}(p+1)}\right)\chi(p+1)p^{n+1}q\beta_{k,\chi}$$

$$\equiv q\sum_{\lambda \bmod p}(1-\bar{\chi}(p+1))\beta_{d,\chi\lambda}\sum_{t=1}^{p-1}\bar{\lambda}(t)t^{d-1}(p^{n-1}(t^{p-1}-1)+p^n t^{p-2}) \pmod{p^{n+2}}, \tag{1.9}$$

where $d \in \{1,2,\dots,p(p-1)\}$ *with* $k \equiv d \pmod{p(p-1)}$.

In [2], Z.-W. Sun proved a Voroni-type congruence for the Euler numbers:

$$\frac{3^{k+1}+1}{4}E_k \equiv \frac{3^k}{2}\sum_{j=0}^{2^n-1}(-1)^{j-1}(2j+1)^k\left\lfloor\frac{3j+1}{2^n}\right\rfloor \pmod{2^n} \tag{1.10}$$

for even $k \geq 0$, where $\lfloor x \rfloor = \max\{z \in \mathbb{Z}: z \leq x\}$. He also showed that the Stern congruence (1.1) is an easy consequence of (1.10). It is not difficulty to check that (1.10) is equivalent to

$$(3^k+1)\beta_{k,\chi_4} \equiv 3^{k-1}\sum_{j=0}^{2^n-1}\chi(2j+1)(2j+1)^{k-1}\left\lfloor\frac{3(2j+1)}{2^{n+1}}\right\rfloor \pmod{2^{n+1}}$$

In this paper, we shall prove the following Voroni-type congruences for $\beta_{n,\chi}$.

Theorem 1.5. *Let* p *be a prime and* a *be an integer with* $p \nmid a$. *Suppose that* χ *is a character modulo* p^m. *Suppose that* $n \geq m$, $k \geq 1$ *and* k *has the same parity as* χ. *If* $p \geq 5$, *or* $p = 2, 3$ *and* $n \geq 2$, *Then*

$$(\chi(a)a^k-1)\beta_{k,\chi} \equiv \chi(a)a^{k-1}\sum_{j=1}^{p^n-1}\chi(j)j^{k-1}\left\lfloor\frac{ja}{p^n}\right\rfloor \pmod{p^n}. \tag{1.11}$$

2. Several auxiliary lemmas

Lemma 2.1. *Suppose that* $m \geq 3$ *and* χ *is a Dirichlet character with the conductor* 2^m. *Then* $\chi(5) \neq 1$ *is a* 2^{m-2}*-th root of unity.*

Proof. Since $a^{2^{m-2}} \equiv 1 \pmod{2^m}$ for any odd a, we have $\chi(5)^{2^{m-2}} = \chi(5^{2^{m-2}}) = 1$. Assume that $\chi(5) = 1$. Note that for every $a \equiv 1 \pmod 4$, there exist $0 \leq j \leq 2^{m-2}-1$ such that $a \equiv 5^j \pmod{2^m}$. Hence $\chi(a) = 1$ for every $a \equiv 1 \pmod 4$. Thus χ can be reduced to a character modulo 4. So we must have $\chi(5) \neq 1$. □

Lemma 2.2. *Suppose that* p *is an odd prime and* χ *is a Dirichlet character with the conductor* p^m. *If* $m > k \geq 1$, *then* $\chi(p^{m-k}+1)$ *is a* p^k*-th primitive root of unity.*

Proof. First, we show that $\chi(p^{m-1}+1)$ is a p-th primitive root of unity. Clearly,

$$\chi(p^{m-1}+1)^p = \chi((p^{m-1}+1)^p) = \chi\left(1+\sum_{j=1}^{p}\binom{p}{j}p^{j(m-1)}\right) = \chi(1) = 1.$$

So we only need to show that $\chi(p^{m-1}+1) \neq 1$. Assume on the contrary that $\chi(p^{m-1}+1) = 1$. Then $\chi(kp^{m-1}+1) = \chi((p^{m-1}+1)^k) = 1$ for each $0 \leq k \leq p-1$. Thus for any $0 \leq k \leq p-1$ and $1 \leq d \leq p^{m-1}-1$ with $p \nmid d$, letting $0 \leq k' \leq p-1$ be the integer such that $dk' \equiv k \pmod p$, we have

$$\chi(kp^{m-1}+d) = \chi(d(k'p^{m-1}+1)) = \chi(d)\chi(k'p^{m-1}+1)) = \chi(d).$$

Hence χ can be reduced to a character modulo p^{m-1}. This leads an contradiction since the conductor of χ is p^m.

Suppose that $k \geq 2$. Similarly, we have $\chi(p^{m-k}+1)$ is a p^k-th primitive root of unity. Note that

$$(p^{m-k}+1)^{p^{k-1}} = 1 + p^{m-1} + \sum_{r=2}^{p^{k-1}} \binom{p^{k-1}-1}{r-1} \frac{p^{m-1+(m-k)(r-1)}}{r} \equiv p^{m-1}+1 \pmod{p^m},$$

since p^{r-1} can't divides r for $r \geq 2$. Then we have $\chi(p^{m-k}+1)^{p^{k-1}} \neq 1$, i.e., $\chi(p^{m-k}+1)$ is a p^k-th root of unity. □

Define

$$S_k(n,\chi) = \sum_{j=1}^{n} \chi(j) j^k.$$

Lemma 2.3. *Suppose that p is a prime and χ is a character modulo p^m, where $m \geq 1$. Then for $n \geq m$ and $k \geq 0$,*

$$S_k(p^n,\chi) \equiv p^{n-m} S_k(p^m,\chi) \pmod{p^n},$$

unless $p=2$, $n=1$ and k is odd.

Proof. We use induction on n. There is nothing to do when $n=m$. Assume that $n>m$ and the assertion holds for smaller values of n. By the induction hypothesis,

$$S_k(p^n,\chi) = \sum_{i=0}^{p-1}\sum_{j=1}^{p^{n-1}} \chi(p^{n-1}i+j)(p^{n-1}i+j)^k \equiv \sum_{i=0}^{p-1}\sum_{j=1}^{p^{n-1}} \chi(j)(kp^{n-1}ij^{k-1}+j^k)$$

$$=pS_k(p^{n-1},\chi) + kp^{n-1}S_{k-1}(p^{n-1},\chi)\sum_{i=0}^{p-1} i \equiv p^{n-m}S_k(p^m,\chi) + \frac{kp^n(p-1)}{2} S_{k-1}(p^{n-1},\chi) \pmod{p^n}.$$

Hence our assertion clearly holds for odd prime p. And when $p=2$, we also have

$$S_{k-1}(2^{n-1},\chi) = \sum_{j=1}^{2^{n-1}} \chi(j) j^{k-1} \equiv \sum_{j=1}^{2^{n-1}} \chi(j) = 0 \pmod{2}.$$

□

Lemma 2.4. *Let p be a prime and χ be a character modulo p^m. Suppose that there exists an integer a with $p \nmid a$ satisfying $\chi(a)a^k - 1$ is prime to p. Then for $n \geq m$,*

$$S_k(p^n,\chi) \equiv 0 \pmod{p^n}.$$

Proof. In view of Lemma 2.3, we need to show that

$$S_k(p^m,\chi) \equiv 0 \pmod{p^m}.$$

Suppose that g is a primitive root modulo p^m and $g^s \equiv a \pmod{p^m}$. Then

$$\begin{aligned} S_k(p^m,\chi) =& \sum_{j=1}^{p^m} \chi(j) j^k \equiv \sum_{i=0}^{p^{m-1}(p-1)-1} \chi(g^i)(g^i)^k = \sum_{i=0}^{p^{m-1}(p-1)-1} (\chi(g)g^k)^i \\ =& \frac{(\chi(g)g^k)^{p^{m-1}(p-1)}-1}{\chi(g)g^k-1} = \frac{g^{kp^{m-1}(p-1)}-1}{\chi(g)g^k-1} \pmod{p^m}. \end{aligned} \tag{2.1}$$

Clearly p^m divides $g^{kp^{m-1}(p-1)}-1$. Assume on the contrary that $\chi(g)g^k-1$ is not prime to p. Then

$$\chi(a)a^k - 1 \equiv \chi(g^s)(g^s)^k - 1 = (\chi(g)g^k)^s - 1 = (\chi(g)g^k-1)\sum_{0\leq i<s} (\chi(g)g^k)^s \pmod{p},$$

i.e., $\chi(a)a^k - 1$ is also not prime to p. So we must have $\chi(g)g^k - 1$ is prime to p, whence p^m divides $S_k(p^m, \chi)$. □

Unfortunately, it is possible that $\chi(g)g^k - 1$ is not prime to p. For example, let χ be the Legendre symbol $\left(\frac{\cdot}{p}\right)$. Then we always have

$$\chi(g)g^{(s+\frac{1}{2})p^n(p-1)} - 1 \equiv 0 \pmod{p^{n+1}}.$$

Lemma 2.5. *Suppose that p is an odd prime and $m \geq 2$. Then for any s with $p \nmid s$, there exists a primitive root g of p^m such that*

$$g^{s(p-1)} \equiv p + 1 \pmod{p^m}.$$

Proof. Let g be an arbitrary primitive root of p^m. Let

$$K = \{1 \leq k \leq p^{m-1}(p-1) : p \nmid k,\ k \equiv 1 \pmod{p-1}\}.$$

Clearly $|K| = p^{m-2}(p-1)$. And for those $k \in K$, g^k are distinct primitive roots modulo p^m. Suppose that

$$g^{ks(p-1)} \equiv a_k p + 1 \pmod{p^m},$$

where $1 \leq a_k \leq p^{m-1}$. Since $p \nmid ks$ and g is also a primitive root modulo p^2, we must have $p \nmid a_k$. Noting that

$$|\{1 \leq a \leq p^{m-1} : p \nmid a\}| = p^{m-2}(p-1),$$

we get

$$\{a_k : k \in K\} = \{1 \leq a \leq p^{m-1} : p \nmid a\},$$

i.e., there exists $k \in K$ such that

$$(g^k)^{s(p-1)} \equiv p + 1 \pmod{p^m}.$$

Then g^k is the desired primitive root. □

In view of Lemma 2.2, $\chi(p+1)$ is a p^{m-1}-th primitive root of unity. So by the definition of d_χ, $\chi(g)$ is a $p^{m-1}d_\chi$-th primitive root of unity. Let $\mathcal{L}$ be the $p^{m-1}d_\chi$-th cyclotomic field. We know

$$(p) = \mathfrak{q}_1^{p^{m-2}(p-1)}\mathfrak{q}_2^{p^{m-2}(p-1)} \cdots \mathfrak{q}_{\phi(d_\chi)}^{p^{m-2}(p-1)},$$

where $\mathfrak{q}_1, \ldots, \mathfrak{q}_{\phi(d_\chi)}$ are distinct prime ideals of $\mathcal{O}_\mathcal{L}$. Recall that $p = \mathfrak{p}_1\mathfrak{p}_2 \cdots \mathfrak{p}_{\phi(d_\chi)}$ in $\mathcal{O}_\mathcal{K}$, where $\mathcal{K}$ is the d_χ-th cyclotomic field. So without loss of generality, we may assume that $\mathfrak{p}_i = \mathfrak{q}_i^{p^{m-2}(p-1)}$ for $1 \leq i \leq \phi(d_\chi)$. On the other hand, since

$$p = (1 - \chi(p+1))^{p^{m-2}(p-1)}a,$$

where a is a unit of the p^{m-1}-th cyclotomic field. Hence

$$(1 - \chi(p+1)) = \mathfrak{q}_1\mathfrak{q}_2 \cdots \mathfrak{q}_{\phi(d_\chi)}.$$

Lemma 2.6. *Suppose that p is an odd prime and χ is a primitive character modulo p^m, where $m \geq 2$. Suppose that g is a primitive root of p^{m+1} such that*

$$g^{p^{m+1}-1} \equiv p + 1 \pmod{p^{m+1}}.$$

Let $\mathfrak{q}$ be a prime ideal factor $\mathfrak{p}$ of p in $\mathcal{O}_\mathcal{L}$. If $\chi(g)g^k - 1$ is divisible by $\mathfrak{q}$, then

$$\frac{\chi(g)g^k - 1}{\bar{\chi}(p+1) - 1} \equiv 1 + \frac{kp}{\chi(p+1) - 1} - \frac{k(k+1)p^2}{2(\chi(p+1) - 1)} \pmod{\mathfrak{q}^{3p^{m-2}(p-1)-1}}.$$

Proof. Without of loss generality, we may assume that $\mathfrak{p}=\mathfrak{q}^{p^{m-2}(p-1)}$ where $\mathfrak{q}$ is a prime ideal factor of p in $\mathcal{O}_{\mathcal{L}}$. Note that for $0\le j\le p^{m+1}$ and $0\le s\le m+1$,

$$\binom{p^{m+1}}{j}=\frac{p^{m+1}}{j}\binom{p^{m+1}-1}{j-1}\equiv 0 \pmod{p^s},$$

unless $p^{m+2-s}\mid j$. So for each $1\le i\le s$, we have

$$\begin{aligned}(\bar{\chi}(g)g^{-k})^{p^{m+1}-1}-1=&\frac{(\bar{\chi}(g)g^{-k}-1+1)^{p^{m+1}}}{\bar{\chi}(g)g^{-k}}-1\\ =&\chi(g)g^k\sum_{j=0}^{p^{m+1}}\binom{p^{m+1}}{j}(\bar{\chi}(g)g^{-k}-1)^j-1\equiv\chi(g)g^k-1 \pmod{\mathfrak{q}^{3p^{m-2}(p-1)}},\end{aligned}$$

by noting that $p^{m-1}\ge p^{m-2}(p-1)+1$ and $\bar{\chi}(g)g^{-k}-1=\bar{\chi}(g)g^{-k}(1-\chi(g)g^k)$ is divisible by $\mathfrak{q}$. On the other hand,

$$\begin{aligned}\bar{\chi}(p+1)-1\equiv&\bar{\chi}(p+1)\bigg((p+1)^{-k}+kp-\frac{k(k+1)}{2}p^2\bigg)-1\\ \equiv&(\bar{\chi}(g)g^{-k})^{p^{m+1}-1}-1+kp\bar{\chi}(p+1)-\frac{k(k+1)}{2}p^2\bar{\chi}(p+1) \pmod{p^3}.\end{aligned}$$

Hence

$$\chi(g)g^k-1\equiv\bar{\chi}(p+1)-1-kp\bar{\chi}(p+1)+\frac{k(k+1)}{2}p^2\bar{\chi}(p+1) \pmod{\mathfrak{q}^{3p^{m-2}(p-1)}}.$$

In particular,

$$\chi(g)g^k-1\equiv\bar{\chi}(p+1)-1 \pmod{\mathfrak{q}^2}.$$

Since those $\mathfrak{q}$ divides $\bar{\chi}(p+1)-1$ but $\mathfrak{q}^2$ doesn't divide $\bar{\chi}(p+1)-1$, we must have $\mathfrak{q}^2$ doesn't divide $\chi(g)g^k-1$. Thus we get

$$\frac{\chi(g)g^k-1}{\bar{\chi}(p+1)-1}=1+\frac{\chi(g)g^k-\bar{\chi}(p+1)}{\bar{\chi}(p+1)-1}\equiv1-\frac{kp\bar{\chi}(p+1)}{\bar{\chi}(p+1)-1}+\frac{k(k+1)p^2\bar{\chi}(p+1)}{2(\bar{\chi}(p+1)-1)} \pmod{\mathfrak{q}^{3p^{m-2}(p-1)-1}}.$$

□

By the proof of Lemma 2.6 and (2.1), we get that for $m\ge2$ and a primitive character χ modulo p^m,

$$(\chi(p+1)-1)S_k(p^m,\chi)\equiv0 \pmod{p^m}. \tag{2.2}$$

Lemma 2.7. *Suppose that $k\ge0$ and χ is a character with the conductor 2^m. If $m\ge3$, or $m=2$ and k is even, or $m=1$ and k is odd, then for $n\ge\max\{m,2\}$,*

$$S_k(2^n,\chi)\equiv0 \pmod{2^n}.$$

Proof. For $m\ge2$, in view of Lemma 2.3, it suffices to prove that

$$S_k(2^n,\chi)\equiv0 \pmod{2^n}.$$

If $m=2$ and k is even, then

$$S_k(2^2,\chi)=1+\chi(3)3^k\equiv1+(-1)\cdot(-1)^k=0 \pmod 4.$$

Assume that $m \geq 3$. We have

$$\begin{aligned} S_k(2^m,\chi) = \sum_{j=1}^{2^m} \chi(j) j^k \equiv \sum_{i=0}^{1} \sum_{j=0}^{2^{m-2}-1} \chi((-1)^i 5^j)(-1)^{ik} 5^{jk} \\ = \Bigl(\sum_{i=0}^{1} \chi(-1)^i (-1)^{ik} \Bigr) \Bigl(\sum_{j=0}^{2^{m-2}-1} \chi(5)^j 5^{jk} \Bigr) = (1+\chi(-1)(-1)^k) \cdot \frac{5^{2^{m-2}k}-1}{\chi(5)5^k-1} \pmod{2^m}. \end{aligned}$$

If $\chi(-1) = (-1)^{k-1}$, there is nothing to do. Suppose that $\chi(-1) = (-1)^k$. Clearly

$$5^{2^{m-2}k} - 1 \equiv 0 \pmod{2^m}.$$

By Lemma 2.1, $\chi(5) \neq 1$ is a 2^{m-2}-th root of unity. And

$$\chi(5)5^k - 1 \equiv \chi(5) - 1 \pmod 4.$$

So $\chi(5)5^k - 1$ is not divisible by $(\chi(5)-1)^2$, i.e., $2(\chi(5)5^k-1)^{-1}$ is 2-integral. So we get that 2^m divides $S_k(2^m,\chi)$.

Finally, when $m = 1$ and k is odd, by Lemma 2.3, we have

$$S_k(2^n,\chi) \equiv 2^{n-2} S_k(2^2,\chi) = 2^{n-2}(1+3^k) \equiv 0 \pmod{2^n}.$$

□

Thus combining Lemmas 2.3, 2.7 and (2.2), we obtain that

Lemma 2.8. *Suppose that p is a prime and χ is a character modulo p^m, where $m \geq 1$. Then for $n \geq m$ and $k \geq 0$,*

$$S_k(p^n,\chi) \equiv 0 \pmod{p^{n-1}}. \tag{2.3}$$

Lemma 2.9. *For an odd prime p and an integer a with $p \nmid a$,*

$$a^{p^n(p-1)q} - 1 \equiv p^{n-m} q (a^{p^m(p-1)} - 1) \pmod{p^{n+m+2}} \tag{2.4}$$

provided that $n \geq m \geq 1$.

Proof.

$$\begin{aligned} a^{p^n(p-1)q} - 1 = (a^{p^m(p-1)} - 1 + 1)^{p^{n-m}q} - 1 = & \sum_{j=1}^{p^n q} \binom{p^{n-m}q}{j} (a^{p^m(p-1)} - 1)^j \\ \equiv & p^{n-m} q (a^{p^m(p-1)} - 1) \pmod{p^{n+m+2}}. \end{aligned}$$

□

3. Proof of Theorem 1.5

For $1 \leq j < p^n$, we write

$$ja = p^n \left\lfloor \frac{ja}{p^n} \right\rfloor + r_j,$$

where $0 \leq r_j < p^n$. For a positive integer k, define $\nu_p(k) = \max\{\alpha \in \mathbb{N} : p^\alpha \mid k\}$. Then

$$\begin{aligned}
&\chi(a)a^k S_k(p^n,\chi) = \sum_{j=1}^{p^n-1} \chi(ja)(ja)^k = \sum_{j=1}^{p^n-1} \chi(ja)\left(p^n \left\lfloor \frac{ja}{p^n} \right\rfloor + r_j\right)^k \\
\equiv& \sum_{j=1}^{p^n-1} \chi(r_j) \sum_{t=0}^{k} \binom{k}{t} p^{tn} \left\lfloor \frac{ja}{p^n} \right\rfloor^t r_j^{k-t} \equiv \sum_{j=1}^{p^n-1} \chi(r_j)\left(r_j^k + \sum_{t=1}^{k} \frac{k}{t}\binom{k-1}{t-1} p^{tn} \left\lfloor \frac{ja}{p^n} \right\rfloor^t r_j^{k-t}\right) \\
\equiv& \sum_{j=1}^{p^n-1} \chi(j)j^k + kp^n \sum_{j=1}^{p^n-1} \chi(ja) \left\lfloor \frac{ja}{p^n} \right\rfloor (ja)^{k-1} + \frac{k(k-1)}{2} p^{2n} \sum_{j=1}^{p^n-1} \chi(ja) \left\lfloor \frac{ja}{p^n} \right\rfloor^2 (ja)^{k-2} \pmod{p^{2n+\nu_p(k)}},
\end{aligned}$$

by noting that $\nu_p(t) \leq t-2$ for $t \geq 3$. Thus

$$\frac{\chi(a)a^k-1}{p^n k} S_k(p^n,\chi) \equiv \chi(a)a^{k-1} \sum_{j=1}^{p^n-1} \chi(j)j^{k-1} \left\lfloor \frac{ja}{p^n} \right\rfloor + \frac{k-1}{2} p^n \sum_{j=1}^{p^n-1} \chi(ja) \left\lfloor \frac{ja}{p^n} \right\rfloor^2 (ja)^{k-2} \pmod{p^n}. \tag{3.1}$$

In particular, if p is odd, then

$$\frac{\chi(a)a^k-1}{p^n k} S_k(p^n,\chi) \equiv \chi(a)a^{k-1} \sum_{j=1}^{p^n-1} \chi(j)j^{k-1} \left\lfloor \frac{ja}{p^n} \right\rfloor \pmod{p^n}. \tag{3.2}$$

On the other hand, we know

$$\begin{aligned}
p^n B_{k,\chi} =& S_k(p^n,\chi) + \sum_{t=1}^{k} \binom{k}{t} p^{tn} B_t S_{k-t}(p^n,\chi) \\
=& S_k(p^n,\chi) - \frac{kp^n}{2} S_{k-1}(p^n,\chi) + \frac{k(k-1)p^{2n}}{12} S_{k-2}(p^n,\chi) + \sum_{t=4}^{k} \frac{k}{t}\binom{k-1}{t-1} p^{tn} B_t S_{k-t}(p^n,\chi).
\end{aligned} \tag{3.3}$$

In view of Lemma 2.8, when $p \geq 5$, or $p = 2,3$ and $n \geq 2$, we have

$$p^n B_{k,\chi} \equiv S_k(p^n,\chi) - \frac{kp^n}{2} S_{k-1}(p^n,\chi) + \frac{k(k-1)p^{2n}}{12} S_{k-2}(p^n,\chi) \pmod{p^{2n+\nu_p(k)}}, \tag{3.4}$$

since pB_t is p-integral by the von Staudt-Clausen theorem.

(i) Suppose that $p \geq 5$, or $p = 3$ and $n \geq 2$. By Lemma 2.8 and (3.4), when $p \geq 5$, or $p = 3$ and $n \geq 2$, we have

$$p^n B_{k,\chi} \equiv S_k(p^n,\chi) - \frac{kp^n}{2} S_{k-1}(p^n,\chi) \pmod{p^{2n+\nu_p(k)}},$$

i.e.,

$$\frac{B_{k,\chi}}{k} \equiv \frac{S_k(p^n,\chi)}{p^n k} - \frac{S_{k-1}(p^n,\chi)}{2} \pmod{p^n}, \tag{3.5}$$

Now by Lemma 2.4,

$$S_{k-1}(p^n,\chi) \equiv 0 \pmod{p^n},$$

since

$$\chi(-1)\cdot(-1)^{k-1} - 1 = -2 \not\equiv 0 \pmod{p}.$$

Thus from (3.2) and (3.5), it follows that

$$(\chi(a)a^k-1)\frac{B_{k,\chi}}{k}\equiv\frac{(\chi(a)a^k-1)}{p^nk}S_k(p^n,\chi)\equiv\chi(a)a^{k-1}\sum_{j=1}^{p^n-1}\chi(j)j^{k-1}\left\lfloor\frac{ja}{p^n}\right\rfloor \pmod{p^n}.$$

(ii) Now let $p=2$. In view of (3.4),

$$2^nB_{k,\chi}\equiv S_k(2^n,\chi)-\frac{k2^n}{2}S_{k-1}(2^n,\chi)+\frac{k(k-1)2^{2n}}{12}S_{k-2}(2^n,\chi) \pmod{2^{2n+\nu_2(k)}}.$$

Furthermore, by (3.1), we have

$$\begin{aligned}&\frac{\chi(a)a^k-1}{k}S_k(2^n,\chi)\\ \equiv&2^n\sum_{j=1}^{2^n-1}\chi(ja)\left\lfloor\frac{ja}{2^n}\right\rfloor(ja)^{k-1}+(k-1)2^{2n-1}\sum_{j=1}^{2^n-1}\chi(ja)\left\lfloor\frac{ja}{2^n}\right\rfloor^2(ja)^{k-2} \pmod{2^{2n}}\end{aligned}$$

and

$$(\chi(a)a^{k-1}-1)S_{k-1}(2^n,\chi)\equiv 2^n(k-1)\sum_{j=1}^{2^n-1}\chi(ja)\left\lfloor\frac{ja}{2^n}\right\rfloor(ja)^{k-2} \pmod{2^{n+1}}.$$

Thus

$$\begin{aligned}\frac{\chi(a)a^k-1}{2^{2n-1}k}S_k(2^n,\chi)-\frac{1}{2^{n-1}}\sum_{j=1}^{2^n-1}\chi(ja)\left\lfloor\frac{ja}{2^n}\right\rfloor(ja)^{k-1}\equiv(k-1)\sum_{j=1}^{2^n-1}\chi(ja)\left\lfloor\frac{ja}{2^n}\right\rfloor^2(ja)^{k-2}\\ \equiv(k-1)\sum_{j=1}^{2^n-1}\chi(ja)\left\lfloor\frac{ja}{2^n}\right\rfloor(ja)^{k-2}\equiv\frac{\chi(a)a^{k-1}-1}{2^n}S_{k-1}(2^n,\chi) \pmod 2.\end{aligned}$$

By Lemma 2.7,

$$S_{k-1}(2^n,\chi)\equiv 0 \pmod{2^n}.$$

Note that

$$\chi(a)a^{k-1}-1\equiv\chi(a)a^k-1 \pmod 2.$$

So

$$\frac{\chi(a)a^{k-1}-1}{2^n}S_{k-1}(2^n,\chi)\equiv\frac{\chi(a)a^k-1}{2^n}S_{k-1}(2^n,\chi) \pmod 2.$$

Hence

$$\frac{\chi(a)a^k-1}{2^nk}S_k(2^n,\chi)-\frac{\chi(a)a^k-1}{2}S_{k-1}(2^n,\chi)\equiv\sum_{j=1}^{2^n-1}\chi(ja)\left\lfloor\frac{ja}{2^n}\right\rfloor(ja)^{k-1} \pmod{2^n}.$$

Furthermore, we have

$$(k-1)S_{k-2}(2^n,\chi)\equiv 0 \pmod{2^n}\equiv 0 \pmod 4.$$

Therefore

$$\begin{aligned}(\chi(a)a^k-1)\frac{B_{k,\chi}}{k}\equiv&\frac{\chi(a)a^k-1}{2^nk}S_k(2^n,\chi)-\frac{\chi(a)a^k-1}{2}S_{k-1}(2^n,\chi)\\ \equiv&\chi(a)a^{k-1}\sum_{j=1}^{2^n-1}\chi(j)j^{k-1}\left\lfloor\frac{ja}{2^n}\right\rfloor \pmod{2^n}.\end{aligned}$$

□

Now applying Theorem 1.5, we have

Theorem 3.1. *Let p be a prime and χ be a character modulo p^m. For any $a \in \mathbb{Z}$ with $p \nmid a$, $(\chi(a)a^k - 1)\beta_{k,\chi}$ is an algebraic integer.*

Proof. By a result of Washington, we know that

$$\begin{aligned}\beta_{n,\chi} &= -\frac{\tau(\chi,\xi_{p^m})}{p^m}\sum_{a=1}^{M-1}\sum_{i=1}^{m}\frac{\bar{\chi}(a)}{i(\xi_{p^m}^a-1)^i}\sum_{j=1}^{i}\binom{i}{j}(-1)^{i-j}j^n\\ &= -\frac{\tau(\chi,\xi_{p^m})}{p^m}\sum_{a=1}^{M-1}\sum_{i=1}^{m}\frac{\bar{\chi}(a)}{(\xi_{p^m}^a-1)^i}\sum_{j=1}^{i}\binom{i-1}{j-1}(-1)^{i-j}j^{n-1},\end{aligned}$$

where ξ_{p^m} is a p^m-th primitive root of unity and

$$\tau(\chi,\xi_{p^m}) = \sum_{a=1}^{p^m}\chi(a)\xi_{p^m}^a.$$

So $\beta_{n,\chi}$ is q-integral for every prime $q \neq p$. However, by Theorem 1.5, $(\chi(a)a^k - 1)\beta_{k,\chi}$ is also p-integral. Thus we must have $(\chi(a)a^k - 1)\beta_{k,\chi}$ is an algebraic integer. □

4. Proof of Theorems 1.1 and 1.2

Since $m \geq 3$, $\chi(5) \neq 1$ is a 2^{m-2}-th primitive root of unity. So $\chi(5) - 1$ divides 2 and

$$\chi(5)5^k - 1 \equiv \chi(5) - 1 \pmod 4.$$

By Theorem 3.1, $(1 - \chi(5))\beta_k$ is an algebraic integer.

Clearly $(2j+1)^{2^nq} \equiv 1 \pmod{2^{n+2}}$. In fact, we have

$$\begin{aligned}(2j+1)^{2^nq} = \sum_{t=0}^{2^nq}\frac{2^nq}{t}\binom{2^nq-1}{t-1}(2j)^t &\equiv 1 + 2^{n+1}qj + 2^{n+1}q(2^nq-1)j^2 + 2^{n+2}q\binom{2^nq-1}{3}j^4\\ &\equiv \begin{cases}1 + 2^{n+1}(qj + 2 - q) & \pmod{2^{n+3}}, \quad \text{if } j \text{ is odd},\\ 1 + 2^{n+1}j & \pmod{2^{n+3}}, \quad \text{if } j \text{ is even}.\end{cases}\end{aligned} \tag{4.1}$$

Let $n_* = \max\{n, m-3\}$ and let a be an integer such that

$$5a \equiv 1 \pmod{2^{n_*+4}}.$$

Since $n_* + 3 \geq m$, with the help of Theorem 1.5,

$$\begin{aligned}&\frac{\chi(a)a^{k+2^nq}-1}{\chi(a)a^{k+2^nq-1}}\cdot\frac{B_{k+2^nq,\chi}}{k+2^nq} - \frac{\chi(a)a^k-1}{\chi(a)a^{k-1}}\cdot\frac{B_{k,\chi}}{k}\\ \equiv&\sum_{j=0}^{2^{n_*+2}-1}\chi(2j+1)(2j+1)^{k+2^nq-1}\left\lfloor\frac{(2j+1)a}{2^{n_*+3}}\right\rfloor - \sum_{j=0}^{2^{n_*+2}-1}\chi(2j+1)(2j+1)^{k-1}\left\lfloor\frac{(2j+1)a}{2^{n_*+3}}\right\rfloor \pmod{2^{n_*+3}}\\ \equiv&2^{n+1}\sum_{\substack{0\leq j<2^{n_*+2}\\ j \text{ is even}}}\chi(2j+1)j\left\lfloor\frac{(2j+1)a}{2^{n_*+3}}\right\rfloor + 2^{n+1}\sum_{\substack{0\leq j<2^{n_*+2}\\ j \text{ is odd}}}\chi(2j+1)(qj+2-q)\left\lfloor\frac{(2j+1)a}{2^{n_*+3}}\right\rfloor \pmod{2^{n+3}}.\end{aligned} \tag{4.2}$$

Furthermore,

$$\begin{aligned}&\frac{(\chi(a)a^{k+2^nq}-1)B_{k+2^nq,\chi}}{\chi(a)a^{k+2^nq-1}(k+2^nq)}-\frac{(\chi(a)a^k-1)B_{k,\chi}}{\chi(a)a^{k-1}k}\\ =&a(1-\bar{\chi}(a)a^{-k-2^nq})\beta_{k+2^nq,\chi}-a(1-\bar{\chi}(a)a^{-k})\beta_{k,\chi}\\ \equiv&a(1-\chi(5)5^{k+2^nq})(\beta_{k+2^nq,\chi}-\beta_{k,\chi})+\chi(5)5^{k-1}(1-5^{2^nq})\beta_{k,\chi}\\ \equiv&5^{-1}(1-\chi(5)5^{k+2^nq})(\beta_{k+2^nq,\chi}-\beta_{k,\chi})-\chi(5)5^{k-1}2^{n+2}(q+2)\beta_{k,\chi}\pmod{2^{n+3}},\end{aligned}\tag{4.3}$$

since

$$5^{2^nq}-1=(4+1)^{2^nq}-1\equiv 2^{n+2}(q+2)\pmod{2^{n+4}}.$$

Thus by (4.2) and (4.3), we get

$$\begin{aligned}&(1-\chi(5))(\beta_{k+2^nq,\chi}-\beta_{k,\chi})-\frac{1-\chi(5)}{1-\chi(5)5^{k+2^nq}}\cdot\chi(5)5^k2^{n+2}(q+2)\beta_{k,\chi}\\ \equiv&\frac{(1-\chi(5))2^{n+1}}{(1-\chi(5)5^{k+2^nq})a}\Bigg(\sum_{\substack{0\le j<2^{n_*+2}\\ j\text{ is even}}}\chi(2j+1)j\left\lfloor\frac{(2j+1)a}{2^{n_*+3}}\right\rfloor+\sum_{\substack{0\le j<2^{n_*+2}\\ j\text{ is odd}}}\chi(2j+1)(qj+2-q)\left\lfloor\frac{(2j+1)a}{2^{n_*+3}}\right\rfloor\Bigg)\\ \equiv&\frac{2^{n+1}}{a}\Bigg(\sum_{\substack{0\le j<2^{n_*+2}\\ j\text{ is even}}}\chi(2j+1)j\left\lfloor\frac{(2j+1)a}{2^{n_*+3}}\right\rfloor+\sum_{\substack{0\le j<2^{n_*+2}\\ j\text{ is odd}}}\chi(2j+1)(qj+2-q)\left\lfloor\frac{(2j+1)a}{2^{n_*+3}}\right\rfloor\Bigg)\pmod{2^{n+3}},\end{aligned}\tag{4.4}$$

since

$$\frac{1-\chi(5)}{1-\chi(5)5^{k+2^nq}}\equiv\frac{1-\chi(5)}{1-\chi(5)}=1\pmod 2.$$

So

$$(1-\chi(5))(\beta_{k+2^nq,\chi}-\beta_{k,\chi})\equiv\frac{1-\chi(5)}{1-\chi(5)5^{k+2^nq}}\cdot\chi(5)5^k2^{n+2}(q+2)\beta_{k,\chi}\equiv\chi(5)2^{n+2}\beta_{k,\chi}\pmod{2^{n+2}}.$$

In particular,

$$(1-\chi(5))\beta_{k+2^nq,\chi}\equiv(1-\chi(5))\beta_{k,\chi}\pmod{2^{n+1}}.$$

Also, we have

$$2\beta_{k,\chi}\equiv 2\beta_{d_2,\chi}\pmod 2\equiv 2\beta_{d_4,\chi}\pmod 4,$$

where $d_i\in\{1,2,\dots,i\}$ satisfying $k\equiv d_i \pmod i$.

Furthermore,

$$\begin{aligned}&\sum_{\substack{0\le j<2^{n_*+2}\\ j\text{ is even}}}\chi(2j+1)j\left\lfloor\frac{(2j+1)a}{2^{n_*+3}}\right\rfloor+\sum_{\substack{0\le j<2^{n_*+2}\\ j\text{ is odd}}}\chi(2j+1)(qj+2-q)\left\lfloor\frac{(2j+1)a}{2^{n_*+3}}\right\rfloor\\ =&\frac14\sum_{j=0}^{2^{n_*+3}-1}\chi(j)(j-1)(1+\chi_4(j))\left\lfloor\frac{ja}{2^{n_*+3}}\right\rfloor+\frac14\sum_{j=0}^{2^{n_*+3}-1}\chi(j)(qj-3q+4)(1-\chi_4(j))\left\lfloor\frac{ja}{2^{n_*+3}}\right\rfloor.\end{aligned}\tag{4.5}$$

By Theorem 1.5,

$$\sum_{j=0}^{2^{n_*+3}-1}\chi(j)j\left\lfloor\frac{ja}{2^{n_*+3}}\right\rfloor\equiv\frac{\chi(a)a^2-1}{\chi(a)a}\beta_{2,\chi}\pmod{2^{n_*+3}},$$

and

$$\sum_{j=0}^{2^{n_*+3}-1}\chi(j)\left\lfloor\frac{ja}{2^{n_*+3}}\right\rfloor\equiv\frac{\chi(a)a-1}{\chi(a)}\beta_{1,\chi}\pmod{2^{n_*+3}}.$$

Thus we have

$$\begin{aligned}&(1-\chi(5))(\beta_{k+2^nq,\chi}-\beta_{k,\chi})-\frac{1-\chi(5)}{1-\chi(5)5^{k+2^nq}}\cdot\chi(5)5^k2^{n+2}(q+2)\beta_{k,\chi}\\ \equiv&2^{n-1}\big((1-\chi(5)5^2)((1+q)\beta_{2,\chi}+(1-q)\beta_{2,\chi\chi_4})+(1-\chi(5)5)(3(1-q)\beta_{1,\chi}+(3q-5)\beta_{1,\chi\chi_4})\big)\pmod{2^{n+3}},\end{aligned}$$

by noting that $\chi\chi_4$ is also a primitive character when $m\geq 3$. Finally, if k and χ are both even, then

$$\frac{(1-\chi(5))5^k}{1-\chi(5)5^{k+2^nq}}\equiv 1\pmod 4$$

and $\beta_{2,\chi\chi_4}=\beta_{1,\chi}=0$. Also, if k and χ are both odd, then

$$\frac{(1-\chi(5))5^k}{1-\chi(5)5^{k+2^nq}}\equiv\frac{(1-\chi(5))5}{1-\chi(5)5}\pmod 4$$

and $\beta_{2,\chi}=\beta_{1,\chi\chi_4}=0$. All are done. □

5. Proof of Theorems 1.3 and 1.4

Proof of the first part of Theorem 1.4. Since $\chi(a)a^k-1$ is prime to $\mathfrak{p}$, in view of Theorem 1.5, both $\beta_{k,\chi}$ and $\beta_{k+p^n(p-1)q,\chi}$ are $\mathfrak{p}$-integral. Let b be an integer such that

$$ab\equiv 1\pmod{p^{n_*+2}}.$$

For $s\geq 0$ and $1\leq t\leq p-1$, by Lemma 2.9,

$$\begin{aligned}(ps+t)^{p^n(p-1)q}=&\sum_{i=0}^{p^nq}\binom{p^nq}{i}(ps)^it^{p^n(p-1)q}\\ \equiv&t^{p^n(p-1)q}+t^{p^n(p-1)q-1}p^{n+1}q\equiv 1+p^nq(t^{p-1}-1)+p^{n+1}qt^{p-2}\pmod{p^{n+2}}.\end{aligned}$$

Thus by Theorem 1.5, we have

$$\begin{aligned}&\frac{\chi(b)b^{k+p^n(p-1)q}-1}{\chi(b)b^{k+p^n(p-1)q-1}}\cdot\beta_{k+p^n(p-1)q,\chi}-\frac{\chi(b)b^k-1}{\chi(b)b^{k-1}}\cdot\beta_{k,\chi}\\ \equiv&\sum_{j=1}^{p^{n_*+2}-1}\chi(j)j^{k-1}(j^{p^n(p-1)q}-1)\left\lfloor\frac{jb}{p^{n_*+2}}\right\rfloor\pmod{p^{n_*+2}}\\ \equiv&\sum_{s=0}^{p^{n_*+1}-1}\sum_{t=1}^{p-1}\chi(ps+t)(ps+t)^{k-1}((ps+t)^{p^n(p-1)q}-1)\left\lfloor\frac{(ps+t)b}{p^{n_*+2}}\right\rfloor\\ \equiv&\sum_{s=0}^{p^{n_*+1}-1}\sum_{t=1}^{p-1}\chi(ps+t)t^{d-1}q(p^n(t^{p-1}-1)+p^{n+1}t^{p-2})\left\lfloor\frac{(ps+t)b}{p^{n_*+2}}\right\rfloor\pmod{p^{n+2}}.\end{aligned}\tag{5.1}$$

Now

$$\begin{aligned}&(1-\bar{\chi}(b)b^{-k-p^n(p-1)q})\beta_{k+p^n(p-1)q,\chi}-(1-\bar{\chi}(b)b^{-k})\beta_{k,\chi}\\ =&(1-\bar{\chi}(b)b^{-k-p^n(p-1)q})(\beta_{k+p^n(p-1)q,\chi}-\beta_{k,\chi})+\bar{\chi}(b)b^{-k}(1-b^{-p^n(p-1)q})\beta_{k,\chi}\\ \equiv&(1-\chi(a)a^{k+p^n(p-1)q})(\beta_{k+p^n(p-1)q,\chi}-\beta_{k,\chi})+\chi(a)a^k(1-a^{p^n(p-1)q})\beta_{k,\chi}\\ \equiv&(1-\chi(a)a^{k+p^n(p-1)q})(\beta_{k+p^n(p-1)q,\chi}-\beta_{k,\chi})+\chi(a)a^kqp^n(1-a^{p-1})\beta_{k,\chi}\pmod{p^{n+1}\mathfrak{p}},\end{aligned}\tag{5.2}$$

where we use Lemma 2.9 in the last inequality. In particular, by (5.1) and (5.2), we have

$$(1-\chi(a)a^{k+p^n(p-1)q})(\beta_{k+p^n(p-1)q,\chi}-\beta_{k,\chi}) \equiv \chi(a)a^k qp^n(a^{p-1}-1)\beta_{k,\chi} \equiv 0 \pmod{p^n\mathfrak{p}}, \tag{5.3}$$

i.e.,

$$\beta_{k+p^n(p-1)q,\chi} \equiv \beta_{k,\chi} \equiv \beta_{d,\chi} \pmod{p^n\mathfrak{p}}.$$

Furthermore,

$$\begin{aligned}&\sum_{s=0}^{p^{n_*+1}-1}\sum_{t=1}^{p-1}\chi(ps+t)t^{d-1}q(p^n(t^{p-1}-1)+p^{n+1}t^{p-2})\left\lfloor\frac{(ps+t)b}{p^{n_*+2}}\right\rfloor\\ =&\sum_{t=1}^{p-1}t^{d-1}q(p^{n-1}(t^{p-1}-1)+p^nt^{p-2})\sum_{\lambda \bmod p}\bar{\lambda}(t)\sum_{j=1}^{p^{n_*+2}-1}\chi\lambda(j)j^{d-1}\left\lfloor\frac{jb}{p^{n_*+2}}\right\rfloor.\end{aligned}$$

By Theorem 1.5,

$$\sum_{j=1}^{p^{n_*+2}-1}\chi\lambda(j)j^{d-1}\left\lfloor\frac{jb}{p^{n_*+2}}\right\rfloor \equiv b(1-\bar{\chi}\bar{\lambda}(b)b^{-d})\beta_{d,\chi\lambda} \equiv \frac{1-\chi\lambda(a)a^d}{a}\beta_{d,\chi\lambda} \pmod{p^{n_*+2}}.$$

Hence

$$\begin{aligned}&\beta_{k+p^n(p-1)q,\chi}-\beta_{k,\chi}-\frac{p^n(a^{p-1}-1)}{1-\chi(a)a^d}\cdot\chi(a)a^dq\beta_{d,\chi}\\ \equiv&\frac{a}{1-\chi(a)a^{k+p^n(p-1)q}}\sum_{t=1}^{p-1}t^{k-1}q(p^{n-1}(t^{p-1}-1)+p^nt^{p-2})\sum_{\lambda \bmod p}\bar{\lambda}(t)\cdot\frac{1-\chi\lambda(a)a^d}{a}\beta_{d,\chi\lambda}\\ \equiv&q\sum_{\lambda \bmod p}\frac{1-\chi\lambda(a)a^d}{1-\chi(a)a^d}\beta_{d,\chi\lambda}\sum_{t=1}^{p-1}\bar{\lambda}(t)t^{d-1}(p^{n-1}(t^{p-1}-1)+p^nt^{p-2}) \pmod{p^{n+1}\mathfrak{p}}.\end{aligned}$$

□

Clearly, the first part of Theorem 1.3 easily follows from (5.3), since $\chi(a)a^k-1$ is prime to any prime ideal factor of p now.

Proof of the second part of Theorem 1.4. Let g be a primitive root modulo p^{m+1} such that

$$g^{p^{m+1}-1} \equiv p+1 \pmod{p^{m+1}}.$$

Let $n_*=\max\{n,m-2\}$ and b be an integer such that

$$gb \equiv 1 \pmod{p^{n_*+3}}.$$

Assume that $\mathfrak{p}=\mathfrak{q}^{p^{m-2}(p-1)}$, where $\mathfrak{q}$ is a prime ideal factor of p in $\mathcal{O}_{\mathcal{L}}$. By Theorem 1.5 and Lemma 2.6, $(1-\bar{\chi}(p+1))\beta_{k,\chi}$ is $\mathfrak{q}$-integral. Similarly as the above discussions, we have

$$\begin{aligned}&(1-\chi(p+1)(p+1)^{k+p^n(p-1)q})(\beta_{k+p^n(p-1)q,\chi}-\beta_{k,\chi})+\chi(p+1)(p+1)^k(1-(p+1)^{p^n(p-1)q})\beta_{k,\chi}\\ \equiv&(p+1)\sum_{s=0}^{p^{n_*+1}-1}\sum_{t=1}^{p-1}\chi(ps+t)t^{d-1}q(p^n(t^{p-1}-1)+p^{n+1}t^{p-2})\left\lfloor\frac{(ps+t)b}{p^{n_*+2}}\right\rfloor \pmod{p^{n+2}},\end{aligned}$$

i.e.,

$$\begin{aligned}&(1-\chi(p+1))(\beta_{k+p^n(p-1)q,\chi}-\beta_{k,\chi})+\frac{(1-\chi(p+1))\chi(p+1)(p+1)^k(1-(p+1)^{p^n(p-1)q})\beta_{k,\chi}}{1-\chi(p+1)(p+1)^{k+p^n(p-1)q}}\\ \equiv&\frac{1-\chi(p+1)}{1-\chi(p+1)(p+1)^{k+p^n(p-1)q}}\sum_{s=0}^{p^{n_*+1}-1}\sum_{t=1}^{p-1}\chi(ps+t)t^{d-1}q(p^n(t^{p-1}-1)+p^{n+1}t^{p-2})\left\lfloor\frac{(ps+t)b}{p^{n_*+2}}\right\rfloor\\ &\pmod{p^{n+2}}.\end{aligned}$$

Now

$$1-\chi(p+1)(p+1)^{k+p^n(p-1)q}\equiv 1-\chi(p+1)(1+dp)\pmod{p^2}$$

So

$$\begin{aligned}\frac{1-\chi(p+1)}{1-\chi(p+1)(p+1)^{k+p^n(p-1)q}}\equiv&\frac{1}{1-\frac{dp\chi(p+1)}{1-\chi(p+1)}}\\ \equiv&1-\frac{dp\chi(p+1)}{\chi(p+1)-1}+\frac{d^2p^2\chi(p+1)^2}{(\chi(p+1)-1)^2}\pmod{p(1-\chi(p+1))}.\end{aligned}$$

On the other hand, with help of Lemma 2.9, we have

$$\begin{aligned}(p+1)^{p^n(p-1)q}-1\equiv&p^{n-1}q((p+1)^{p(p-1)}-1)\\ \equiv&p^{n-1}q\bigg(\binom{p(p-1)}{1}p+\binom{p(p-1)}{2}p^2+\binom{p(p-1)}{3}p^3\bigg)\\ \equiv&p^{n+1}q\bigg(-1+\frac{3}{2}p-\frac{1}{3}p^2\bigg)\pmod{p^{n+3}}.\end{aligned}$$

Now

$$\begin{aligned}&\frac{1-\chi(p+1)}{1-\chi(p+1)(p+1)^{k+p^n(p-1)q}}\cdot\chi(p+1)(p+1)^k((p+1)^{p^n(p-1)q}-1)\beta_{k,\chi}\\ \equiv&\bigg(1-\frac{dp\chi(p+1)}{\chi(p+1)-1}+\frac{d^2p^2\chi(p+1)^2}{(\chi(p+1)-1)^2}\bigg)\cdot\chi(p+1)(1+dp)p^{n+1}q\bigg(-1+\frac{3}{2}p-\frac{1}{3}p^2\bigg)\beta_{k,\chi}\\ \equiv&\bigg(-1+\frac{3}{2}p-dp-\frac{p^2}{3}+\frac{dp\chi(p+1)}{\chi(p+1)-1}-\frac{d^2p^2\chi(p+1)^2}{(\chi(p+1)-1)^2}\bigg)\chi(p+1)p^{n+1}q\beta_{k,\chi}\pmod{p^{n+2}},\end{aligned}$$

by noting that $(1-\chi(p+1))\beta_{k,\chi}$ is p-integral. Furthermore,

$$\begin{aligned}&(p+1)\sum_{s=0}^{p^{n_*+1}-1}\sum_{t=1}^{p-1}\chi(ps+t)t^{d-1}(p^n(t^{p-1}-1)+p^{n+1}t^{p-2})\left\lfloor\frac{(ps+t)b}{p^{n_*+2}}\right\rfloor\\ \equiv&\sum_{\lambda\bmod p}(1-\chi\lambda(p+1)(p+1)^d)\beta_{d,\chi\lambda}\sum_{t=1}^{p-1}\bar{\lambda}(t)t^{d-1}(p^{n-1}(t^{p-1}-1)+p^nt^{p-2})\pmod{p^{n+2}}\\ \equiv&\sum_{\lambda\bmod p}\bigg(1-\chi(p+1)\bigg(1+dp+\frac{d(d-1)}{2}p^2\bigg)\bigg)\beta_{d,\chi\lambda}\sum_{t=1}^{p-1}\bar{\lambda}(t)t^{d-1}(p^{n-1}(t^{p-1}-1)+p^nt^{p-2})\\ &\pmod{p^{n+2}}.\end{aligned}$$

Note that

$$\begin{aligned}
&\sum_{\lambda \bmod p}(1-\chi(p+1))\beta_{d,\chi\lambda}\sum_{t=1}^{p-1}\bar{\lambda}(t)t^{d-1}((t^{p-1}-1)+pt^{p-2})\\
\equiv&\sum_{\lambda \bmod p}(1-\chi(p+1)(p+1)^{-d})\beta_{d,\chi\lambda}\sum_{t=1}^{p-1}\bar{\lambda}(t)t^{d-1}((t^{p-1}-1)+pt^{p-2})\\
\equiv&\frac{1}{p+1}\sum_{s=0}^{p^{n_*+1}-1}\sum_{t=1}^{p-1}\chi(ps+t)t^{d-1}(p(t^{p-1}-1)+p^2t^{p-2})\left\lfloor\frac{(ps+t)(p+1)}{p^{n_*+2}}\right\rfloor\\
\equiv&0 \pmod{p(1-\chi(p+1))}.
\end{aligned}$$

Hence

$$\begin{aligned}
&(1-\chi(p+1))(\beta_{k+p^n(p-1)q,\chi}-\beta_{k,\chi})\\
&-\left(-1+\frac{3}{2}p-dp-\frac{p^2}{3}+\frac{dp\chi(p+1)}{\chi(p+1)-1}-\frac{d^2p^2\chi(p+1)^2}{(\chi(p+1)-1)^2}\right)\chi(p+1)p^{n+1}q\beta_{k,\chi}\\
\equiv&q\left(1-\frac{dp\chi(p+1)}{\chi(p+1)-1}\right)\sum_{\lambda \bmod p}\left(1-\chi(p+1)\left(1+dp+\frac{d(d-1)}{2}p^2\right)\right)\cdot\beta_{d,\chi\lambda}\cdot\\
&\cdot\sum_{t=1}^{p-1}\bar{\lambda}(t)t^{d-1}(p^{n-1}(t^{p-1}-1)+p^nt^{p-2})\\
\equiv&q\sum_{\lambda \bmod p}(1-\chi(p+1))\beta_{d,\chi\lambda}\sum_{t=1}^{p-1}\bar{\lambda}(t)t^{d-1}(p^{n-1}(t^{p-1}-1)+p^nt^{p-2}) \pmod{p^{n+2}}.
\end{aligned}$$

Clearly the terms $\frac{d^2p^2}{(\chi(p+1)-1)^2}$ will vanish unless $p=3$ and $m=2$.

□

Acknowledgment. We are grateful to Professors Zhi-Hong Sun and Zhi-Wei Sun for their helpful discussions on Stein's congruence.

REFERENCES

[1] Z.H.Sun, *Euler numbers modulo* $2n$, Bull. Aust. Math. Soc., **82**(2010), 221-231 .

[2] Z. W. Sun, *On Euler numbers modulo powers of two*, J. Number Theory, **115** (2005), 371-380.

[3] S. S. Wagstaff, Jr., *Prime divisors of the Bernoulli and Euler numbers*, in: Number Theory for the Millennium, III (Urbana, IL, 2000), 357C374, A K Peters, Natick, MA, 2002.

DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING 210093, PEOPLE'S REPUBLIC OF CHINA
E-mail address: `haopan79@yahoo.com.cn`

DEPARTMENT OF BASIC COURSE, NANJING INSTITUTE OF TECHNOLOGY, NANJING 211167, PEOPLE'S REPUBLIC OF CHINA
E-mail address: `yongzhang1982@163.com`